Cakota | Environmental Quality


Agricultural Groundwater Monitoring Program

Pipestem Creek Aquifer


Foster, Stutsman, and Wells Counties

Aquifer At-a-Glance				
Area	23.5 square miles			
Aquifer Type	Unconfined Surficial			
Major Land Uses over Aquifer	Crops (55%)			
(percentage of aquifer area covered in 2017) ¹	Grassland/Pasture (31%)			
Depth to Water* 2	5-12 feet			
Total Unique Wells Sampled	7			
Wells Sampled in 2016	2			
Samples Collected in 2016	4			
Years Sampled	1996, 2001, 2006, 2011, 2016			
*Depths to water may vary seasonally, year to year, and across the aquifer				

- Aguifer materials consist of sands and gravels that were
- deposited in an ancient valley by streams moving meltwater away from glaciers during the last ice age.^{2,3}
- The aquifer averages about 20 feet thick. 2,3
- Domestic, irrigation, and stock wells are installed in the aquifer.
- In North Dakota, permits are required to withdraw large quantities of groundwater. In 2019, 325 gallons of permitted water were drawn from the aquifer; irrigation use consumed the largest quantity of water. For more information on water use and permits, contact the North Dakota State Water Commission (swc.nd.gov).



2019 Pipestem Creek aquifer permitted water use (from North Dakota State Water Commission (swc.nd.gov)) ↓



About the Agricultural Groundwater Monitoring Program

- The North Dakota Department of Environmental Quality monitors a network of wells in approximately 50 surficial aquifers that are at elevated risk of agricultural contamination.
- Aquifers are sampled on a 5-year rotation.
- Monitoring began in 1992.
- The vast majority of these aquifers are located in central and eastern North Dakota.
- Water is tested for 21 general chemistry parameters, eight trace metals, and 64 pesticides.

References

- US Department of Agriculture, 2017, National Agricultural Statistics Service Cropland Data Layer.
 Buturla, F. Jr., 1970, Geology and Ground Water Resources of Wells County, North Dakota, North Dakota State Water Commission County Ground Water Studies 12-Part 3, North Dakota Geological Survey Bulletin 51.
- (3) Trapp, H. Jr., 1968, Geology and Ground Water Resources of Eddy and Foster Counties, North Dakota, North Dakota State Water Commission County Ground Water Studies 5-Part 3, North Dakota Geological Survey Bulletin 44.


Water Chemistry

Is Aquifer Water High in...?


	Analyte	Result	2016 Median	Potential Effects
	Analyte	Nesuit	Concentration	i otentiai Enects
	Arsenic	Locally	< 0.005 mg/L	Skin or circulatory system damage, increased cancer risk
r	Iron	YES	0.76 mg/L	Metallic taste/odor, discoloration of surfaces
	Manganese	YES	0.42 mg/L	
?	Sodium	YES	106 mg/L	Taste, people with certain health conditions may need to limit intake
	Sulfate	YES	262 mg/L	Taste/odor, laxative effect for people not used to the water
	For more information of out Manipurus Contention at Local (MOLs), booking officers and the observation of outliness contentions for the contention of the content outliness outl			

For more information about Maximum Contaminant Levels (MCLs), health effects, and treatment options for these contaminants and more, see the NDDEQ's fact sheets (deq.nd.gov/wq/1_Groundwater) or visit the US EPA website (epa.gov/ground-water-and-drinking-water).


Dominant Water Type	Water Hardness
Calcium-Bicarbonate	Very Hard







Pesticides



State Pesticide Management Plan

Agricultural Groundwater Monitoring Program aquifers are monitored as a part of the State Pesticide Management Plan. A Prevention Action Level (PAL) threshold of 25% of the pesticide's Maximum Contaminant Level (MCL)* or Health Advisory Level (HAL) is used to identify whether action is needed to prevent further contamination.

Prevention Action Level Exceedances	None
MCL or HAL Exceedances	None

Number of Unique Wells with Pesticide Detections since 1996

1 of 7 Total Wells

2016 Pesticide Detections				
Picloram	1 Well	Herbicide applied to crops and roads/rights-of-way		